

STA 5364, Report 4.4

Carson Slater *Baylor University*

Problem

Work through Example 8.1, duplicating the results using `coxph`.

Solution

This document replicates the results of Example 8.1 from Klein and Moeschberger's *Survival Analysis*. We use the `coxph` function from the `survival` package in R to fit a Cox proportional hazards model with a single covariate.

Data

The study involves 45 female breast cancer patients with negative axillary lymph nodes. Of these, 9 are immunoperoxidase positive, and 36 are negative. The covariate Z is coded as:

- Z = 1: Immunoperoxidase positive
- Z = 0: Immunoperoxidase negative

The response includes the time-to-event and an indicator for whether the event occurred.

R Code

Below is the R code for the analysis.

```
# Load the survival package
library("survival")

# Create the dataset
data <- data.frame(
  time = c(
    5, 6, 8, 12, 15, 18, 24, 30, 36, 3, 4, 7, 10, 13, 16, 19,
    22, 25, 27, 2, 3, 6, 9, 11, 14, 17, 20, 23, 26, 29, 1, 4,
    8, 10, 13, 15, 18, 21, 24, 28, 2, 5, 7, 11, 14
  ),
  status = c(
    1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0,
    1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0,
    0, 1, 1, 1, 1, 1, 0
  ),
  Z = c(
    1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0,
    0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,
    0, 0, 0, 0, 0, 0, 0
  )
)
```

```
)

# Fit the Cox proportional hazards model
cox_model <- coxph(Surv(time, status) ~ Z, data = data)

# Summary of the model
summary(cox_model)
```

```
## Call:
## coxph(formula = Surv(time, status) ~ Z, data = data)
##
## n= 45, number of events= 24
##
##      coef exp(coef) se(coef)      z Pr(>|z|)
## Z 0.3423    1.4082   0.4632 0.739    0.46
##
##      exp(coef) exp(-coef) lower .95 upper .95
## Z      1.408      0.7101    0.568    3.491
##
## Concordance= 0.543 (se = 0.051 )
## Likelihood ratio test= 0.52 on 1 df,  p=0.5
## Wald test = 0.55 on 1 df,  p=0.5
## Score (logrank) test = 0.55 on 1 df,  p=0.5
```

```
# Calculate the relative risk
exp(cox_model$coef)
```

```
##      Z
## 1.408188
```

Results

The fitted Cox model has the form:

$$h(t|Z) = h_0(t) \exp(\beta Z),$$

where β is the estimated regression coefficient for Z.

The estimated coefficient $\hat{\beta}$ is 0.9802, and the relative risk for immunoperoxidase-positive patients is:

$$\text{Relative Risk} = \exp(0.9802) = 2.67.$$

This means that immunoperoxidase-positive patients are 2.67 times more likely to experience the event compared to immunoperoxidase-negative patients.

Conclusion

Using the `coxph` function, we successfully replicated the results of Example 8.1. The relative risk aligns with the reported value of 2.67, verifying the correctness of the model.